



## Science Drivers for Multiwavelength Investigations Using the New Gamma-Ray Observatories and Missions

Dave Thompson  
GLAST LAT Multiwavelength Coordinator

### Outline

1. The Multiwavelength Revolution
2. Topics of Special Interest to the Gamma-ray Community

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 1



## A Revolution? Why?

Simple literature search using NASA ADS.  
Search for the word "Multiwavelength" in titles.

1976	0
1986	2
1996	58
2006	97

Many astrophysicists have come to recognize  
the value of multiwavelength research.

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 2



## One Reason – Facilities

The situation across the spectrum in 1982 – 25 yrs ago:

Radio telescopes – VLA open 2 years, no VLBA

Sub-mm – just getting started

IR – no satellites (IRAS 1983), detectors limited

Optical – largest telescopes: Hale 5 m, Soviet 6 m, few  
CCDs

UV – IUE (a bright spot)

X-ray – Hakucho Japanese satellite

Gamma-ray – COS-B turned off in that year

TeV – Whipple Observatory just getting started

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 3



## Another Reason – Communication

1977 – 111 computers on ARPANET, forerunner of the  
Internet. Most information exchange was on paper.

1981 – First IBM PC

1983 – TCP/IP protocol; SENDMAIL program started;  
e-mail became practical.

1988 – NSFNET backbone upgraded to 1.5 Mbps

1990 – Tim Berners-Lee and the first Web server

1993 – Mosaic Web browser

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 4




The major scientific driver for multiwavelength research comes from the fact that we mostly call ourselves astrophysicists, not astronomers. We want to explore what is there, but also to understand how those things work.

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 5



## An Example – The Vela Pulsar



D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 6





## Multiwavelength Research: The Gamma-ray Perspective

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 7



### Gamma-ray Facilities: More Numerous, More Capable




D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 8



## Known Gamma-ray Sources Are Multiwavelength



Gamma-ray sources are nonthermal, typically produced by interactions of high-energy particles.

Known classes of gamma-ray sources are multiwavelength objects, seen across much of the spectrum.

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 9



## What Do Gamma-ray Measurements Offer?

- Huge energy range – 9+ orders of magnitude
- All-sky coverage, from both ground and space
- Excellent sensitivity compared to previous instruments
- Good source locations – 1 arcmin in many cases
- High time resolution for individual photons
- Imaging for some extended sources

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 10



## Some Other Needs for Astrophysics

- Distance – redshift, Dispersion Measure, proper motion, column density
- Composition – spectroscopy
- Precise source locations and imaging
- Velocities
- Polarization
- Magnetic fields
- Theories to connect the observations to physical models



D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 11




## Source Identification: What Is It?

First Clue: Gamma-ray variability Radio sources in the error box




One flat-spectrum radio source, 260 mJy at 5 GHz; one marginally-flat source, 49 mJy; other sources are much weaker



Optical observations:

The 49 mJy source is a normal galaxy;  
The 260 mJy source has an optical counterpart with a redshift  $z=0.83$



Spectral energy distribution is bimodal like other blazars  
Conclusion: a flat spectrum radio quasar (FSRQ) –

Wallace et al., 2002


Variable optical polarization is seen.  
Only an X-ray upper limit found.

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 12



## Time-Dependent Modeling: How Does It Work?




D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 13



## Incomplete Measurements: What Are the Implications?

RX J1713.7-3946 – Is This SNR Accelerating Cosmic Ray Protons?



Lemoine-Goumard, et al. 2007

D. Thompson

Multiwavelength Workshop Adler Planetarium Chicago Aug. 9-10, 2007 14



## Summary

Multiwavelength studies are valuable to many aspects of astrophysics.

They are particularly important to maximize the scientific return from the new gamma-ray facilities.